

RESEARCH ARTICLE

IMMUNOSTIMULANT AND FREE RADICAL SCAVENGING STUDIES OF *GANODERMA APPLANATUM*

Madhu. Divakar^{1*}, Lins Mary Joy²

¹PPG College of Pharmacy, Coimbatore, Tamilnadu, India

²Nirmala College of Pharmacy, Muvattupuzha, Ernakulum, Kerala, India

Received: 17 Dec, 2022/Revision: 03 Jan, 2023 /Accepted: 11 Feb, 2023

Abstract: Objective: The present animal experimentation study is planned to screen the immunostimulant efficacy, free radical scavenging potential and the acute toxicity of the bracket fungus, *Ganoderma applanatum*. **Methods:** The study analyse the invitro immunostimulant activity by phagocytic index determination using *Candida albicans*, acute toxicity study by brine shrimp assay and the free radical scavenging activities of *G. applanatum* solvent extracts. **Conclusion:** The percentage immunostimulation was found to be 88% for the chloroform-methanol (1:1) *Ganoderma applanatum* extract (GAE). The free radical scavenging activity studies were performed by utilising invitro model of hydroxyl, superoxide and lipid peroxide radical generating system. **Results** indicated that GAE at 100 mcg/ml showed 88.1, 89.19, and 81.87 % scavenging activity of hydroxyl, superoxide and lipid peroxide radical respectively. The LC50 was determined using brine shrimp assay method and calculated as 875 mcg /ml. The present study shows that *G. applanatum* can be utilised as an effective natural supplement to impart immunostimulation.

KEYWORDS: *Ganoderma applanatum*, Immunostimulant activity, free radical scavenging activity

INTRODUCTION:

In Chinese traditional medicine, *G. applanatum* has been used commonly as haemostatic, immunostimulant, tumour inhibitor, and also for the treatment of rheumatic tuberculosis and oesophageal carcinoma^{1,2}. The common synonyms are artist's bracket, bear bread, artist's conk etc³. *G. applanatum* is a parasitic and saprophytic fungus lives inside the living or dead tree wood as mycelium. Compounds

like applanoxidic acid and sugars like arabinol, ribose, fucose, mannitol, sorbitol, glucose, sucrose, maltose, uronic acid etc were isolated and reported previously^{5,6,7}. Mohammad S.H *et al* reported⁸ the usefulness of *G. applanatum* in the management of diabetes meletus, hyperlipidemia etc. The present study is designed to evaluate the invitro immunostimulant activity by phagocytic index determination using *Candida albicans*, acute toxicity study by *brine shrimp assay* and the free radical scavenging activities of GAE.

Corresponding Author:
Dr Madhu.C.Divakar,
PPG College of Pharmacy, Coimbatore, Tamilnadu, India
Email- madhu.divakar@gmail.com



MATERIALS AND METHODS:

The dried and matured fruiting bodies of *Ganoderma applanatum* (Ganodermataceae) were collected from Agriculture University, Trivandrum, Kerala in January 2003. The specimen was identified at the Dept. of Plant Pathology, Agriculture University, Vellayani, Trivandrum. A voucher specimen deposited at the herbarium of the department of Pharmacognosy, SRIPMS, Coimbatore -641044, India.

The 'chloroform: methanol soxhleted extract of *G. applanatum* (yield: 8.0%w/w) shows positive reactions for steroids and triterpenoids upon phytochemical screening⁴.

Studied activities

Immunostimulant activity was conducted by phagocytic index determination (Table-4, Fig-1) and free radical scavenging activity studies (for hydroxyl, superoxide and lipid peroxide radicals) by invitro studies. The LC₅₀ determination was performed by Brine shrimp assay (BSA) method¹⁰.

Acute toxicity studies: Brine Shrimp Assay method

Brine shrimp assay method¹⁰ was followed to find out the LC₅₀ for the extract GAE. The brine shrimp eggs were hatched in a rectangular chamber filled with artificial sea water and five numbers each were transferred to vials using a 9-inch disposable pipette. The survival rate of the shrimps was observed after 24 h for different concentrations of GAE. The LC₅₀ was found from the dose -response graph. The results are tabulated in Table 1.

Table -1. Toxicity studies

No.	Con: (mcg/ml)	% Inhibition	LC ₅₀ mcg/ml
1	250	0	875
2	500	20	
3	750	40	
4	1000	60	

Free radical scavenging activity studies

• Hydroxyl radical scavenging activity¹¹

This study was conducted by measuring the inhibition of deoxyribose degradation in presence of the test drug extracts. Hydroxyl radical was generated by Fe EDTA and H₂O₂ in presence of ascorbic acid. The extract GAE was added in various concentrations. (10mcg/ml – 100mcg/ml) to a reaction mixture containing deoxyribose (3mM), FeCl₃ (20mM/pH 7.4) to make a final volume of 3ml. To this mixture, trichloroacetic acid and thiobarbituric acid (0.5ml each) were added and measured the absorbance at 532nm. The percentage of hydroxyl radical inhibition and IC₅₀ of the test drug extracts were determined by the method of Halliwell et. al (1987). A 0.01 mM copper sulphate solution was used as a reference standard. The results are tabulated in Table 2.

Table-2. Hydroxyl radical scavenging activity

No.	Con: (mcg/ml)	% Inhibition	EC ₅₀ mcg/ml
1	10	10.04±0.5185	56
2	30	30.43±0.08576	
3	50	45.08±0.0989	
4	70	62.34±0.01416	
5	100	88.179±0.0476	
6	CuSO ₄	95.6±0.05185	

• Superoxide radical scavenging activity^{12,14}

Superoxide radical scavenging activity was studied according to the method reported in the literature. Alkaline dimethyl sulphoxide (1% in 5mM NaOH) was added to the reaction mixture containing nitro-blue tetrazolium (NBT 0.1mg) and the test drug GAE at various concentrations. The absorbance was determined at 560nm. The reduction of NBT by the superoxide radical generated was calculated in the presence and absence of test drugs. In this study, thio urea (20mM) was used as the reference standard. The results were tabulated in Table 3.

Table -3. Superoxide radical scavenging activity

No.	Con: (mcg/ml)	% Inhibition	EC ₅₀ mcg/ml
1	10	17.37	28
2	30	35.59	
3	50	53.38	
4	70	70.33	
5	100	89.19	
6	Thiourea	90.04	

• Lipid peroxide scavenging activity

In this study, the liver tissue homogenate¹³ of albino rats was prepared in phosphate buffer saline of pH 7.4. The protein content of the homogenate was adjusted to 10mg/ml. The effect of the test compounds on lipid peroxide was estimated as malondialdehyde by thiobarbituric acid (TBA) method. To the reaction mixture containing test drug extracts at various concentrations, 1ml of liver tissue homogenate and 1ml of "HCl thiobarbituric acid-trichloro acetic acid reagent" was added. The mixture was warmed gently for 5min in a water bath at 37°C. After cooling the flocculent precipitate was removed by centrifugation at 1000rpm for 10min. The absorbance of the supernatant liquid was measured at 532nm against blank and the lipid peroxide content was determined using the extinction coefficient $1.56 \times 10^5 \text{m}^{-1}\text{cm}^{-1}$. The final result was expressed as nanomoles of malondialdehyde per mg of protein. Vitamin E (50mcg/ml) was used as the standard reference in this study. The results are tabulated in Table 4.

Table-4. Lipid peroxide scavenging activity

No.	Con: (mcg/ml)	% Inhibition	EC ₅₀ mcg/ml
1	10	16.1	51
2	30	32.55	
3	50	50.16	
4	70	65.43	
5	100	81.87	
6	Vit E	88.08	

• Immunostimulant activity studies⁹

The immunostimulant activity study was conducted by phagocytic index determination using *Candida albicans*. Human blood (2-3 drops) was taken by finger prick method from a healthy volunteer with prior informed consent and documented with the institutional ethics committee (IEC/SRIPMS/11/18). The slide with the blood collected was kept on a cotton pad in a sterile petridish and incubated at 37°C for 25 min. After incubation the clot was removed very gently and the slide was slowly drained with sterile normal saline taking care not to wash the adhered neutrophils. The slide was flooded with predetermined concentration of the test drug, incubated at 37°C for 15 min and flooded with a suspension of *Candida albicans* in Hank's balanced salt solution and human serum and incubated at 37°C for 1h. After this, the slide was drained, fixed with methanol and stained with Giesma stain. The mean number of phagocytised cells on the slide was determined microscopically for 100 granulocytes. This number was taken as the phagocytic index (PI) and was compared with the basal phagocytic index of control. The results are tabulated in Table 5. The percentage immunostimulation was calculated by using the equation.

$$\% \text{ immunostimulation} = \frac{\text{PI}_{(T)} - \text{PI}_{(C)}}{\text{PI}_{(C)}} \times 100$$

PI_(T) – Phagocytic index of test
PI_(C) – Phagocytic index of control

Table-5. Immunostimulant activity studies

Drug Groups (n)	Dose mg/ml	Phagocytic index	% Immunostimulation
GAE	50	41 ± 0.816	64 ± 3.26
GAE	100	47 ± 0.47	88 ± 0.87**
Control (normal saline)	-	25 ± 0.8164	-
n=5 p value < 0.01**, < 0.05* GAE: <i>Ganoderma applanatum</i> extract			




Figure 1. Phagocytosis of WBC with *Candida albicans*

CONCLUSION:

The qualitative analysis of GAE showed the presence of steroidal triterpenes. The extract showed significant scavenging of superoxide, hydroxyl, and lipid peroxide radicals, when compared to the standards CuSO_4 , Vit.E, and Thiourea respectively. The study showed that the extract obtained from *Ganoderma applanatum* (GAE) can be used as an effective natural immunostimulant and free radical scavenging agent with less toxicity for the affected individuals in the post Covid-19 pandemic period.

REFERENCES:

- [1]. Vijay H.M., Comtois P., Sharma R., Lemieux R. Allergenic components of *Ganoderma applanatum*. Grana. 1991; 30(1): 167–170.
- [2]. Ginns, James. Polypores of British Columbia (Fungi: Basidiomycota). Victoria, BC: Province of British Columbia - Forests, Lands, and NR Operations; 2017. p. 105. ISBN 978-0-7726-7053-3.
- [3]. Goldman, Gary B. Field Guide to Mushrooms & Other Fungi in South Africa. Marieka Gryzenhout. Cape Town, South Africa; 2019. ISBN 978-1-77584-654-3. OCLC 1117322804.
- [4]. IL Finar. Stereochemistry and Chemistry of Natural Products, Volume 2, 5th edn. 1975; p. 158.
- [5]. Boldizsar I, Horvath K, Szedlay K.G, Molnar-Perl. Simultaneous GC-MS quantitation of acids and sugars in the hydrolyzates of immunostimulant, water-soluble polysaccharides of basidiomycetes. Chromatographia. 1998; 47(7-8): 413–419.
- [6]. Elkhatieb WA, Zaghlol GM, El-Garawani IM, Ahmed EF, Rateb ME, Abdel Moneim AE. *Ganoderma applanatum* secondary metabolites induced apoptosis through different pathways: in vivo and in vitro anticancer studies. Biomedicine & Pharmacotherapy. 2018; 101: 264–277.
- [7]. Moradali MF, Mostafavi H, Hejaroude GA, Tehrani AS, Abbasi M, Ghods S. Investigation of potential antibacterial properties of methanol extracts from fungus *Ganoderma applanatum*. Chemotherapy. 2006; 52(5): 241–244. doi:10.1159/000094866
- [8]. Hossain MS, Barua A, Hoque Tanim MA, Hasan MS, Islam MJ, Hossain MR, Emon NU, Hossen SM. *Ganoderma applanatum* mushroom provides new insights into the management of diabetes mellitus, hyperlipidemia, and hepatic degeneration: A comprehensive analysis. Food Sci Nutr. 2021 Aug; 9(8): 4364–4374.
- [9]. Kulkarni SR, Karnade VS. Study of immunopotentiating activity of naphthoquinone extracts of leaves of *Lawsonia alba*. Indian Drugs. 1983; 35(7): 427–433.
- [10]. Mayer BN, Ferrigni NR, Putname JE, Jacobson LB, Nichols DE, McLaughlin JL. Brine shrimp: A convenient general bioassay for active plant constituents. Planta Med. 1982; 45: 31–34.
- [11]. Halliwell B, Gutteridge JMC, Aruoma OI. The deoxyribose method: A simple test tube assay for determination of rate constants for reactions of hydroxyl radicals. Analytical Biochemistry. 1987; 165: 215–219.
- [12]. Savitha C, Thiripathi P. Indian Journal of Experimental Biology. 1995; 33: 428.
- [13]. Rajshree CR, Rajmohan T, Augusti KT. Antiperoxide effect of garlic protein in alcohol-fed rats. Indian Journal of Experimental Biology. 1998 Jan; 36(1): 60–64.

[14]. Hyland K, Voisin E, Banoun H, Auclair C.
Analytical Biochemistry. 1983; 135: 280–287.

Cite of article: CD Madhu, JM Lins. Immunostimulant and free radical scavenging studies of *Ganoderma applanatum*.
Int. J. Med. Lab. Res. 2023; 8(1): 21–25. <http://doi.org/10.35503/IJMLR.2023.8103>

CONFLICT OF INTEREST: Authors declared no conflict of interest

SOURCE OF FINANCIAL SUPPORT: Nil

International Journal of Medical Laboratory Research (IJMLR) - Open Access Policy

Authors/Contributors are responsible for originality of contents, true references, and ethical issues.

IJMLR publishes all articles under Creative Commons Attribution- Non-Commercial 4.0 International License (CC BY-NC). <https://creativecommons.org/licenses/by-nc/4.0/legalcode>