

CHILDHOOD MALIGNANCIES PATTERN IN KAMPALA, UGANDA

*Ivan Emmanuel, Samuel david, odongo moses, Bwambale Elijah

Kampala Hospital, Central Region, Uganda

Received: 18 June 2016/Revised: 25 June 2016/Accepted: 1 July 2016

Abstract: Childhood cancers represent crucial health problem worldwide. This is especially so in poorer countries, where childhood cancer detected in advance stage, so prognosis became poorer and also appropriate treatment is either not available or not affordable. Many children are never diagnosed at all, many are diagnosed very late, and when a diagnosis is made the treatment options may be limited. Therefore this 10-year retrospective study was undertaken to evaluate and documentation of the pattern of childhood cancers in local region. **Methods:** We reviewed 533 childhood (≤ 15 years) malignancies diagnosed at the cancer department of our hospital in a 10-year period. **Results:** The 533 malignancies comprised 7.8% of all cancers. The malignancies frequently seen in early childhood (1-10 years) accounted for 88%. Retinoblastoma (30.8%), Burkitt lymphoma (17.8%) and acute leukemia (15.9%) were the most common pediatric cancers. **Conclusions:** Childhood cancer with malignant lymphomas and leukemia is the most common. Retinoblastoma was alone is the most prominent type of malignancy. The significant increase in the number of cases and male excess in lymphoid malignancies requires further research into environmental agents that may be implicated.

Key words: acute leukemia; cancer; retinoblastoma; Burkitt lymphoma

INTRODUCTION:

Childhood cancers represent major health problem issue worldwide. Developing countries are suffering more with the childhood cancer because first of all they have the resource problem and then the detection was done on the advance stage so that the prognosis became poorer and also appropriate treatment is either not available or not affordable¹. Children's are diagnosed at very late which make limitation in their treatment options. Pediatric cancer has become an increasingly important cause of childhood mortality and a major public health concern. In view of the higher birth rates and younger population in the developing countries, childhood malignancies are accordingly more prevalent than in the developed world. Childhood cancers comprise just 0.5%-2% of malignancies in the industrialized countries,

but 4.3%-12.5% in the developing countries. Not surprisingly, over 80% of the global childhood cancers are estimated to occur in the developing countries.² In the low income developing countries, pediatric cancer survival is low, largely because of the poor access to good quality pediatric oncology care. Such data are essential for optimal channeling of scarce healthcare resources in the developing countries.^{3,4} The need for up to date cancer data is particularly crucial in view of the striking ethnic and variations in regional that have been documented, sometimes even within a country⁵. Climatic, genetic and other environmental factors often exert considerable influence on the pattern of malignant tumors. Even within the African country differences have been noted in the pattern of childhood malignant tumors⁶. Understanding of the pattern of childhood malignancies in different geographical zones support the hypothesis of environmental factors in the etiopathogenesis of cancers in children.

Address of corresponding author:

Ivan Emmanuel

Kampala Hospital,

Central Region, Uganda

METHODS:

This study include 10-year (2004-2014) retrospective review of all childhood (0-15 years old) malignancies diagnosed at the oncology department of kampala hospital, central region, Uganda. Case files of all children admitted with malignancies in the hospitals during the study period were retrieved and relevant data were extracted and recorded in the study format. Such data included socio-demographic characteristics (age, sex, socio-economic class), types of the malignancy and affected area of body. Patients whose case files were missing or those with incomplete data were excluded from the study.

RESULTS:

A total of 533 childhood (0-15 years) malignancies were diagnosed and analyse for the 10-year study period, which comprised 39% of all pediatric cases and 9.6% of all cancers. 288 male and 245 female patients were reported, which had ratio of 1.2:1. Childhood malignancy were prominently seen in group of 1-5 years and group of 6-10 years, accounting for 48% and 40% respectively comparing with Malignancy in group of >10 years and group of <1 year. Retinoblastoma was the commonest childhood malignancy found in children with 30.8% of all childhood malignancies while Burkitt's lymphoma was the second most frequently encountered childhood malignancy with 17.8%, followed by leukemia (15.9%). From leukemia subtypes, acute lymphoblastic leukemia (ALL) is more commonly found in comparison to acute myeloid leukemia (AML), while other subtypes were relatively less in numbers.

Table 1. Type of malignancy observed

Malignancy	No.	%	Male	Female	<1 y	1-5 y	6-10 y	>10 y
Retinoblastoma	164	30.8	91	73	10	148	20	-
Burkitt's lymphoma	95	17.8	52	43	-	21	49	24
Leukemia	85	15.9	49	36	3	35	40	16
Rhabdomyosarcoma	40	7.5	15	25	-	15	11	11
Nephroblastoma	41	7.69	23	18	4	15	6	-
Other non-Hodgkin lymphoma	23	4.32	14	9	-	2	13	9
Hodgkin lymphoma	18	3.38	12	6	-	2	7	9
Malignant testicular tumors	4	0.75	4	-	-	1	2	-
Malignant ovarian tumors	13	2.44	-	13	-	-	5	7
Neuroblastoma	17	3.19	9	8	-	11	3	2
Skin Cancer	4	0.75	1	3	-	1	1	2
Kaposi sarcoma	7	1.31	5	2	-	2	3	2
Malignant bone tumors	7	1.31	4	3	-	1	2	2
Other sarcomas	15	2.81	9	6	3	2	3	8
Total	533	100	288	245	20	256	165	92

Leukemia most commonly found in the 1-10 year age group, particularly the group of 6-10 years. Other (non- retinoblastoma) malignant embryonal tumors (neuroblastoma, rhabdomyosarcoma) were the 1/4th of the all the cancers, collectively comprising 15.2% of all childhood malignancy. The hemato-lymphoid malignancies (41.4%) (leukemias, lymphomas) were the most prevalent found, followed by embryonal cancers, namely retinoblastoma, nephroblastoma, neuroblastoma and embryonal rhabdomyosarcoma.

DISCUSSION:

The childhood malignancy with 9.6% of all cancers during the study period, higher than developed countries but consistent with studies in other parts of Africa this difference in developed and developing countries because of higher birth rates and which leads to larger proportion of young people and children in their population. Slight male preponderance (M:F=1.2:1) found, which is comparable to other around the world varying from 1.2:1 to 1.5:⁷⁻¹⁰ The peak age incidence of pediatric malignancies was 48% in early childhood (1-5 years), followed by 40% in late childhood (6-10 years). Retinoblastoma which was the commonest childhood cancer found predominately in group of 1-6 years while in developed countries the commonest cancer was acute leukemia.^{11,12} High rates of retinoblastoma have also been documented in other African countries with this ophthalmic tumor among the largest childhood cancers in Tanzania, Ghana, Kenya, Malawi, Zambia, Congo. Cancer treatment is generally expensive and often times requiring prolonged hospital stay. The parents of these patients need to expense the costs of treatment including drugs, diagnostic investigations, meals and hospital stay. Therefore, many families of children with cancer experience financial difficulties. In developed countries medical expenses is paid by their health insurance plan for patients treatment. For individuals without health insurance or who need financial assistance to cover care costs, resources are available, including government sponsored programs and services supported by voluntary organizations¹³.

On the other hand, in resource poor countries where health insurance and re-sources to help families with children with malignancies through financial difficulties are virtually nonexistent. Burkitt lymphoma was the leading childhood cancer, in other African studies and the peak age group was in late childhood (5-9 years)^{8,-10}. In Lagos, retinoblastoma has displaced Burkitt lymphoma as the commonest childhood malignancy¹⁴. Leukemia (mostly acute leukemia), the 3rd most frequent childhood malignancy comprised 15.9%. Though somewhat higher than several other Sub-Saharan African studies,^{14,15} In the United States, Europe, Middle East and the Indian subcontinent, leukemia comprised a third of childhood cancers^{1,17,18}. In addition, unlike in several solid malignancies which often permit survival for several months, acute leukemia can be rapidly fatal, thereby obviating late diagnosis. acute leukemia incidence was more is consistent with the other studies¹⁹. The high incidence of AML has also been reported among black American children in late childhood^{20,21}.

CONCLUSION:

Childhood cancer with malignant lymphomas and leukemia is the most common. Retinoblastoma was alone is the most prominent type of malignancy. The significant increase in the number of cases and male excess in lymphoid malignancies requires further research into environmental agents that may be implicated.

REFERENCES:

1. Paraic R. Childhood cancer: Rising to the challenge. *UICC* 2006; 5:5- 43
2. Barr R, Riberio R, Agarwal B, Masera G, Hesselning P, Magrath I. Pediatric Oncology in Countries with Limited Resources. In: Pizzo PA, Poplack DG, eds. *Principles and Practice of Pediatric Oncology*, 5th ed. Philadelphia: Lippincott Williams and Wilkins, 2006: 1605-1617.
3. Gurney JG, Bondy ML. Epidemiology of childhood cancer. In: Pizzo PA, Poplack DG, eds. *Principles and Practice of Pediatric Oncology*, 5th ed. Philadelphia: Lippincott Williams and Wilkins, 2006: 2-14.
4. Howard SC, Metzger ML, Wilimas JA, Quintana Y, Pui CH, Robison LL, et al. Childhood cancer epidemiology in low-income countries. *Cancer* 2008;112:461-472.
5. Wessels G, Hesselning PB. Unusual distribution of childhood cancer in Namibia. *Pediatr Hematol Oncol* 1996;13:9-20.
6. Mahmoud A. O., Buari M. B., Adekoya B. S. Pattern of orbito ocular growths in Ilorin, Nigeria. *The Tropical Journal of Health Sciences* 2007; 14:23-27.
7. Akinde OR, Abdulkareem FB, Daramola AO, Anunobi CC, Banjo AA. Morphological pattern of childhood solid tumours in Lagos University Teaching Hospital. *Nig Q J Hosp Med* 2009;19:169-174.
8. Samaila MO. Malignant tumours of childhood in Zaria. *Afr J Paediatr Surg* 2009;6:19-23.
9. Akinde OR, Abdulkareem FB, Daramola AO, Anunobi CC, Banjo AA. Morphological pattern of childhood solid tumours in Lagos University Teaching Hospital. *Nig Q J Hosp Med* 2009;19:169-174.
10. Welbeck JE, Hesse AA. Pattern of childhood malignancy in Korle Bu Teaching Hospital. *West Afr J Med* 1998;17:81-84.
11. Michel G, von der Weid NX, Zwahlen M, Adam M, Rebholz CE, Kuehni CE, et al. The Swiss Childhood Cancer Registry: rationale, organisation and results for the years 2001-2005. *Swiss Med Wkly* 2007;137:502-509.
12. Linet MS, Ries LA, Smith MA, Tarone RE, Devesa SS. Cancer surveillance series: recent trends in childhood cancer incidence and mortality in the United States. *J Natl Cancer Inst* 1999;91:1051-1058.
13. The World Bank Report: World Development Indicators. Available from <http://www.data.worldbank.org/news>. Assessed on 14th March, 2012.
14. Tijani SO, Elesha SO, Banjo AA. Morphological pattern of pediatric solid cancer in Lagos, Nigeria. *West Afr J Med* 1995;14:174-180.
15. Mukiibi JM, Banda L, Liomba NG, Sungani FC, Parkin DM. Spectrum of childhood cancers in Malawi 1985-1993. *East Afr Med J* 1995;72:25-29.
16. Patil PS, Elem B, Gwavava NJ, Urban MI. The pattern of pediatric malignancy in Zambia (1980-1989): a hospital-based histopathological study. *J Trop Med Hyg* 1992;95:124-127.
17. Jamal S, Mamoon N, Mushtaq S, Luqman M. Pattern of childhood malignancies: study of 922 cases at Armed Forces Institute of Pathology (AFIP), Rawalpindi, Pakistan. *Asian Pac J Cancer Prev* 2006;7:420-422.
18. Arora RS, Eden TO, Kapoor G. Epidemiology of childhood cancer in India. *Indian J Cancer* 2009;46:264-273.
19. Belson M, Kingsley B, Holmes A. Risk factors for acute leukaemias in children. *Environ Health Perspectives* 2007;115:138-145.
20. Deschler B, Lübbert M. Acute myeloid leukemia: epidemiology and etiology. *Cancer* 2006;107:2099-2107
21. Bhatia S, Neglia JP. Epidemiology of childhood acute myelogenous leukemia. *J Pediatr Hematol Oncol* 1995;17:94-100

Conflict of interest:

The authors declare that there are no areas of conflicting interest.