

## RESEARCH ARTICLE

### DEVELOPMENT OF NEW SOLVENT SYSTEMS FOR THE ANALYSIS OF DIAZEPAM FROM BLOOD

AK Jaiswal<sup>1</sup>, Vinod Kumar<sup>2</sup>, Shiva Prasad<sup>3</sup>

<sup>1</sup>Dept of Forensic Medicine, All India Institute of Medical Sciences, New Delhi-110029,

<sup>2</sup>Institute of Pesticide Formulation Technology, Gurugram -122016, Haryana

<sup>3</sup>Department of Microbiology, Orissa University of Agriculture and Technology, Orissa, Bhubaneswar

Received: 18 April, 2020/ Revision: 21 April, 2020/ Accepted: 20 May, 2020

**ABSTRACT:** Diazepam is a medication of the benzodiazepine family that typically produces a calming effect. It is commonly used to treat a range of conditions including anxiety, alcohol withdrawal syndrome, muscle spasms, seizures, trouble sleeping, and restless legs syndrome. Diazepam was analyzed by using different instrumentation techniques such as HPLC, GLC, LC-MS, GC-MS etc. An attempt has been made to develop a new method for analysis of Diazepam in Biological sample such as blood using High Performance Thin Layer Chromatography (HPTLC) Plate. Diazepam was extracted from blood using liquid-liquid extraction method and analyzed by using High Performance Thin Layer Chromatography Plate. For Chromatographic separation, various binary and tertiary solvent systems were used as mobile phase. Fifteen different solvent systems with different volumetric ratios were used in separation and identification of Diazepam. Developed plates were viewed under UV light followed by spray of chromogenic reagents which successfully increased the sensitivity without dispensing the simplicity of the method. The method developed is a simple, rapid, expensive, non-destructive and reproducible which can be performed in any laboratory easily.

**KEYWORDS:** Diazepam, Benzodiazepine, Solvent Systems, TLC, Rf, Spraying Reagent etc.

### INTRODUCTION:

Diazepam was invented by Dr. Leo Steinbach Hoffmann-La Roche at the company's Nutley, New Jersey, facility following chlordiazepoxide (Librium), which was approved for use in 1960.<sup>[1]</sup> It is commonly used to treat a range of conditions including anxiety, alcohol withdrawal syndrome, muscle spasms, seizures, trouble sleeping, and restless legs syndrome. It may also be used to cause memory loss during certain medical procedures. It can be taken by mouth, inserted into

the rectum, injected into muscle or vein. When given into a vein, effects begin in one to five minutes and last up to an hour. By mouth, effects may take 40 minutes to begin. Common side effects include sleepiness and trouble with coordination. Serious side effects are rare. Decreased breathing, and an increased risk of seizures if used too frequently. Long term use can result in tolerance, dependence,

**Corresponding Author:**

**Dr. A K Jaiswal**

**Department of Forensic Medicine and Toxicology, All India Institute of Medical Sciences, New Delhi-110029**



and withdrawal symptoms on dose reduction. Abrupt stopping after long-term use can be potentially dangerous. After stopping, cognitive problems may persist for six months or longer. It is not recommended during pregnancy or breastfeeding. Its mechanism of action is by increasing the effect of the neurotransmitter *gamma*-Amino butyric acid (GABA). Diazepam was first synthesized by Leo Steinbach, and was first manufactured by Hoffmann-La Roche. Routinely, Diazepam was analysed by using Thin Layer Chromatography (TLC)<sup>[2-4]</sup> High Performance Liquid Chromatography (HPLC)<sup>[5-7]</sup>, UV-Visible Spectrophotometer<sup>[8]</sup>, Gas Liquid Chromatography (GLC)<sup>[9-11]</sup>, Gas Chromatography-Mass Spectroscopy (GC-MS)<sup>[12]</sup> and Liquid Chromatography-Mass Spectroscopy (LC-MS)<sup>[13]</sup>. It has been one of the most frequently prescribed medications in the world since its launch in 1963<sup>[14]</sup>. In the United States it was the highest selling medication between 1968 and 1982, selling more than two billion tablets in 1978 alone<sup>[15]</sup>. In 1985 the patent ended, and there are now more than 500 brands available on the market. Diazepam is on the World Health Organization's List of Essential Medicines, the most effective and safe medicines needed in a health system<sup>[16]</sup>. The wholesale cost in the developing world is about 0.01 USD per dose as of 2014<sup>[17]</sup>. In the United States it is about 0.40 USD per dose<sup>[18]</sup>. Diazepam can be administered orally, intravenously (must be diluted, as it is painful and damaging to veins) intramuscularly (IM), or as a suppository<sup>[19]</sup>.

## **MATERIAL AND REAGENTS:**

### ***Chemicals***

Methanol, Ammonia, Chloroform, Acetic Acid, Toluene, Butanol, Benzene, Diethyl ether, Ethanol, Conc. Hydrochloric acid, Sulphuric acid, anhydrous Sodium sulphate, n-Hexane, Carbon Tetrachloride, Xylene, N,N-Diethylformamide, Sodium tungstate dihydrate and Amyl alcohol, Cyclohexane,

Acetonitrile, Paraffin Paper, Iodine etc from Merck India were used.

### ***Glassware***

Beaker, conical flask, chromatographic chamber, measuring cylinder, separating funnel, volumetric flask, TLC spreyer from Borosil India were used. Microcapillaries from Top Tech Lab Equipments Pvt. Ltd. was used.

### ***Ultrapure water***

Ultrapure water from Rions India was used throughout.

### ***High Performance Thin layer chromatography (HPTLC) plates***

HPTLC Silicagel 60 F<sub>254</sub> aluminium sheets from Merck Germany was used.

### ***Miscellaneous***

UV chamber, oven, Scale, Pencil, Digital weighing balance etc from different manufacturer were used.

#### **• Preparation of Standard solution**

• 1000 ppm solution of Diazepam was prepared by dissolving 0.1 gm of Diazepam in 100 ml of methanol.

#### **• 3.Extraction of Diazepam Poison from Blood**

5ml of blood sample was taken in a beaker, pinch of sodium tungstate and 1-2 ml of H<sub>2</sub>SO<sub>4</sub> was added to it and heated at 60°C for 2-3 minutes. Contents are filtered through filter paper and washed with 0.1N sulphuric acid. The filtrate was made alkaline with ammonia and mixed with ether:chloroform (3:1) and shaken well. Organic layer was collected. The collected organic layer was dried. The dried residue is mixed with few drops of methanol which was used for spotting on TLC plate.

#### **• Spotting of sample and standard on HPTLC plates**

Diazepam extracted from blood was loaded on the HPTLC plates along with the standard.

#### **• Development of the TLC Plates**

Spotted plates were developed in different solvent systems taken in different ratios (Table 1). After

developing, the TLC plates were taken out from the solvent chamber and air-dried.

• **Visualization of TLC Plates**

The dried and developed TLC plates were first observed under UV and then kept in iodine chamber for visualization.

**RESULT AND DISCUSSION:**

The dried and developed TLC plates were first observed under UV and then kept in iodine chamber for visualization. There were 15 different solvent systems (Table 1) with different volumetric ratios were used. The R<sub>f</sub> value was also varying with different volumetric ratios. The R<sub>f</sub> value of Diazepam extracted from blood under experimental conditions was found nearly equal to standard. The response of separation of diazepam in all the 15 solvent system were observed along R<sub>f</sub> value and presented in the Table 1.

**Table 1. Solvent system with ratio and R<sub>f</sub> values**

S.No.	Solvent System	Ratios	R <sub>f</sub> of Standard	R <sub>f</sub> of Sample
1.	Diethyl ether: acetone	5:5	0.97	0.97
		6:4	0.98	0.98
		7:3	0.98	0.98
		8:2	0.92	0.92
		9:1	0.97	0.97
		4:6	0.96	0.96
		3:7	0.97	0.96
		2:8	0.98	0.97
		1:9	0.98	0.98
2.	Chloroform: methanol	5:5	0.97	0.92
		6:4	0.86	0.85
		7:3	0.84	0.83
		8:2	0.86	0.85
		9:1	0.87	0.85
		4:6	0.84	0.82
		3:7	0.89	0.85
3.	Chloroform: Toluene	5:5	0.90	0.88
		6:4	0.93	0.92
		7:3	0.78	0.66
		8:2	0.79	0.77

4.	Diethyl ether: Methanol	9:1	0.88	0.85
		4:6	0.92	0.91
		3:7	0.89	0.85
		2:8	0.92	0.92
		1:9	0.82	0.82
		5:5	0.87	0.84
		6:4	0.89	0.87
		7:3	0.88	0.87
		8:2	0.89	0.86
		9:1	0.86	0.82
		4:6	0.88	0.88
		3:7	0.87	0.87
		2:8	0.89	0.85
		1:9	0.88	0.85
		5.	Ethanol: Methanol	5:5
6:4	0.72			0.71
7:3	0.78			0.74
8:2	0.72			0.71
9:1	0.80			0.78
4:6	0.77			0.76
3:7	0.87			0.86
2:8	0.91			0.85
1:9	0.88			0.85
6.	Diethyl ether: Carbon tetrachloride	5:5	0.25	0.23
		6:4	0.30	0.30
		7:3	0.39	0.36
		8:2	0.39	0.35
		9:1	0.52	0.49
		4:6	0.28	0.28
		3:7	0.29	0.27
		2:8	0.31	0.26
		1:9	0.31	0.28
7.	Acetic acid: Acetonitrile	5:5	0.79	0.78
		6:4	0.91	0.91
		7:3	0.83	0.80
		8:2	0.80	0.78
		9:1	0.77	0.69
		4:6	0.82	0.78
8.	Xylene: Diethyl ether	3:7	0.89	0.82
		2:8	0.83	0.74
		1:9	0.76	0.72
		5:5	0.24	0.24
		6:4	0.34	0.28
		7:3	0.22	0.22
		8:2	0.17	0.17
9:1	0.80	0.60		
4:6	0.32	0.31		
3:7	0.33	0.33		

	2:8	0.41	0.38		7:3	0.86	0.81
	1:9	0.39	0.36		8:2	0.89	0.79
9. Dimethyl	5:5	0.86	0.84		9:1	0.76	0.71
formamide:	6:4	0.86	0.82		4:6	0.78	0.74
Toluene	7:3	0.84	0.80		3:7	0.88	0.82
	8:2	0.89	0.86		2:8	0.91	0.87
	9:1	0.87	0.84		1:9	0.95	0.91
	4:6	0.88	0.85	15. Cyclohexane:	5:5	0.97	0.92
	3:7	0.83	0.82	acetone	6:4	0.88	0.83
	2:8	0.89	0.88		7:3	0.82	0.76
	1:9	0.91	0.89		8:2	0.92	0.85
10 Acetic acid:	5:5	0.83	0.78		9:1	0.78	0.71
Butanol	6:4	0.83	0.83		4:6	0.88	0.88
	7:3	0.79	0.74		3:7	0.86	0.82
	8:2	0.83	0.80		2:8	0.79	0.71
	9:1	0.79	0.75		1:9	0.91	0.89
	4:6	0.87	0.87				
	3:7	0.87	0.87				
	2:8	0.84	0.83				
	1:9	0.81	0.81				
11 Acetic acid:	5:5	0.75	0.72				
Carbon tetra	6:4	0.73	0.71				
chloride	7:3	0.76	0.76				
	8:2	0.79	0.76				
	9:1	0.70	0.66				
	4:6	0.71	0.68				
	3:7	0.68	0.62				
	2:8	0.55	0.48				
	1:9	0.75	0.73				
	5:5	0.70	0.67				
12. Acetic acid:	6:4	0.89	0.85				
Toluene	7:3	0.91	0.89				
	8:2	0.88	0.86				
	9:1	0.95	0.92				
	4:6	0.75	0.73				
	3:7	0.75	0.70				
	2:8	0.52	0.47				
	1:9	0.27	0.23				
	5:5	0.99	0.98				
13. Amyl alcohol:	6:4	0.95	0.93				
cyclohexane	7:3	0.98	0.98				
	8:2	0.55	0.51				
	9:1	0.51	0.47				
	4:6	0.98	0.97				
	3:7	0.74	0.70				
	2:8	0.82	0.78				
	1:9	0.86	0.80				
	5:5	0.89	0.86				
14. Amyl alcohol:	6:4	0.80.	0.76				
Chloroform							

### CONCLUSION:

Method developed for the detection of diazepam in blood is very simple, easy to perform in any laboratory with very less time and reproducible result. This method can be utilized by Forensic Scientist/Chemist of different Forensic Science Laboratory.

### REFERENCE:

- [1] Pollack, Andrew (June 26, 2012). "Roche to Shut Former U.S. Headquarters" *New York Times*. Archived from the original on March 31, 2013. Retrieved January 10, 2014.
- [2] Azzam RM, Notarianni LJ, Ali HM, Rapid and simple chromatographic method for the determination of diazepam and its major metabolites in human plasma and urine. *J Chromatogr B Biomed Sci Appl.* 1998; 708,1-2: 304-309.
- [3] Gjerde H, Dahlin E, Christophersen AS, Clinical impairment of benzodiazepines-relation between benzodiazepine concentrations and impairment in apprehended drivers. *J Pharm Biomed Anal* 1992;10: 317-322.
- [4] Fisher LE, Perch S, Bonfiglio MF, Geers SM, Simultaneous determination of midazolam and flumazenil concentrations in human plasma by gas chromatography. *J.Chromatogr. B: Biomed. Sci. and Appl.* 1995;665: 217-221.

- [5] Hudecova T, Batorova V, Hatrik S, Havranek E, Validation of an HPLC method for the analysis of decomposition products in injectable diazepam. *Ceska Slov Farm.*2004; 53,5:228-33.
- [6] Srikantha D, Raju RR, RP-HPLC Estimation of Imipramine Hydrochloride and Diazepam in Tablets. *Indian J Pharm Sci.*2015; 77,3: 343-347.
- [7] Tulja RG, Gowri SD, Kadgapathi P, Satyanarayana BD, Development of an RP-HPLC method for the simultaneous estimation of propranolol hydrochloride and diazepam in combined dosage form, *Ind J Pharm Edu Res.* 2011; 45,4: 296-300.
- [8] El-Hawary WF, Issa YM, Talat A, Spectrophotometric Determination of Diazepam in pure form, tablets and ampoules. *Int J Biomed Sci.* 2007; 3,1: 50-55.
- [9] De Silva JAF, Schwartz MA, Stefanovic V, Kaplan Arconte LD, Determination of Diazepam (Valium) in Blood by Gas Liquid Chromatography. *Anal. Chem.* 1964; 36,11: 2099-2105.
- [10] Dhar AK, Kutt H, Monitoring Diazepam and Desmethyldiazepam concentrations in plasma by Gas-Liquid Chromatography with use of a nitrogen sensitive detector, *Clin. Chem.*1979; 25,1: 137-140.
- [11] Gamble JAS, Assaf RAE, Mackay JS, Margaret, Kennedy S, Howard PJ, Estimation of plasma diazepam: Critique of a method using gas-liquid chromatography and benzene extraction. *Anaesthesia* 1975; 30: 159-163.
- [12] Goldberger BA, Chronister CW, Merves ML, Quantitation of benzodiazepines in blood and urine using gas chromatography-mass spectrometry (GC-MS). *Methods Mol Biol.*2010; 603: 75-87.
- [13] Stephanie J M, Rebecka C, Miles M, Gwendolyn AMM, Quantitation of Benzodiazepines in urine, serum, plasma and meconium by LC-MS-MS. *Journal of Analytical Toxicology* 2008; 32: 491-498.
- [14] Calcaterra NE, Barrow JC, Classics in Chemical Neuroscience: Diazepam (Valium). *ACS Chem. Neurosci.* 2014; 5, 4: 253-260.
- [15] Nicholas E. Calcaterra and James C. Barrow Classics in Chemical Neuroscience: Diazepam (Valium). *ACS Chem. Neurosci.* 2014; 5,4: 253-260.
- [16] "WHO Model List of Essential Medicines (19th List)"(PDF). *World Health Organization.* 2015.
- [17] "Diazepam". *International Drug Price Indicator Guide.* Archived from the original on 28 March 2017. Retrieved 2<sup>nd</sup> Dec. 2015.
- [18] Diazepam". The American Society of Health-System Pharmacists. Archived from the original on 2015-06-30. Retrieved 2015-06-05.
- [19] Pere Munne (1998). M. Ruse, ed. "Diazepam" Inchem.org.Inchem.org. Archived from the original on 2006-02-27. Retrieved 2006-03-11.

**Cite of article:** Jaiswal AK, Kumar V, Prasad S. Development of new solvent systems for the analysis of diazepam from blood . *Int. J. Med. Lab. Res.* 2020; 5,2:27-31. <http://doi.org/10.35503/IJMLR.2020.5204>

**CONFLICT OF INTEREST:** Authors declared no conflict of interest

**SOURCE OF FINANCIAL SUPPORT:** Nil

- ✓ International Journal of Medical Laboratory Research (IJMLR) - Open Access Policy
- ✓ Authors/Contributors are responsible for originality of contents, true references, and ethical issues.
- ✓ IJMLR publishes all articles under Creative Commons Attribution- Non-Commercial 4.0 International License (CC BY-NC). <https://creativecommons.org/licenses/by-nc/4.0/legalcode>